

2007 Annual Report

Pallid Sturgeon Population Assessment and Associated Fish Community Monitoring for the Missouri River: Segment 1



**Prepared for the U.S. Army Corps of Engineers – Missouri River Recovery Program
By:**

Tyler Haddix, Landon Holte and Cindy Lott

**Montana Fish, Wildlife & Parks
Government Bldg.
East Kansas Street
Fort Peck, MT 59223**

April 2008

EXECUTIVE SUMMARY

Segment 1 of the Missouri River was sampled during the sturgeon and fish community seasons of the Pallid Sturgeon Population Assessment Program (Program) during 2007. This was the second year segment 1 has been sampled according to the Program's Standard Operating Procedures. Segment 1 is used as a reference bend do to it being the most altered river segment from Fort Peck Dam, MT to the headwaters of Lake Sakakawea, ND. Comparisons made between segment 1 and other less altered segments should allow for inferences to how Fort Peck Dam is affecting the Missouri River downstream.

Although pallid sturgeon *Scaphirynchus albus* have been continuously stocked within 4.5 river miles of segment 1 for the last four years, no pallid sturgeon have been sampled in segment 1 during either 2006 or 2007. Similarly, sturgeon chub *Macrhybopsis gelida* and sicklefin chub *M. meeki*, two potential prey have not been sampled in segment 1. Conversely, shovelnose sturgeon *Scaphirynchus platorynchus*, blue suckers *Cycleptus elongatus* and sand shiners *Notropis stramineus* have been collected in both years of sampling. Other native target species that have been collected in one of the two years of sampling include sauger *Sander canadense* and western silvery minnow *Hybognathus argyritis*.

A total of 12,362 fish consisting of 13 species were collected in segment 1 during 2007, with the vast majority (99.7%) being collected in the mini fyke nets set in the fish community season. This was substantially higher than the total number of fish sampled during 2006 (N = 719). Of the 2007 total, 81.8% of were juvenile white suckers and 9.5% fathead minnows. We observed lower CPUE for shovelnose sturgeon in both trammel and otter trawl during both seasons in 2007 compared to 2006. Higher flows out of the Milk River in June of 2007 may have had an influence on the higher relative abundance of catostomids and cyprinids observed in 2007, when compared to 2006, when the Milk River was almost stagnant.

TABLE OF CONTENTS

Introduction.....	1
Study Area	4
Methods.....	5
Sample site selection and description	5
Sampling gear	6
Data Collection and Analysis.....	8
Results	
Pallid sturgeon	11
Targeted Native River Species.....	11
Missouri River Fish Community	20
Discussion.....	21
Acknowledgments.....	23
References.....	24
Appendices.....	25

LIST OF TABLES

Table 1. Presence/absence of all fishes collected in segments 1 through 3 of the Missouri River in 2006 and 2007.....19

LIST OF FIGURES

Figure 1a. Map of segment 1 of the Missouri River with major tributaries, common landmarks, and historic stocking locations for pallid sturgeon. Segment 1 encompasses the Missouri River from Fort Peck Dam (River Mile 1771.5) to the confluence of the Milk River (River Mile 1760).10

Figure 2. Abundance of all fishes sampled in all gears within segment 1 of the Missouri River during 2006 and 2007, by season.....14

Figure 3. Mean annual catch-per-unit-effort by season of all fishes collected using trammel nets in segment 1 of the Missouri River during 2006 and 2007.15

Figure 4. Mean annual catch-per-unit-effort by season of all fishes collected using otter trawls in segment 1 of the Missouri River during 2006 and 2007.16

•
Figure 5. Mean annual catch-per-unit-effort by season of all fishes collected using mini fyke nets in segment 1 of the Missouri River during 2006 and 2007.17

Figure 6. Length frequency histogram and length-weight relationship for all shovelnose sturgeon captured in segment 1 of the Missouri River during 2006 and 2007.18

LIST OF APPENDICES

Appendix A. Phylogenetic list of Missouri River fishes with corresponding letter codes used in the long-term pallid sturgeon and associated fish community sampling program.	26
Appendix B. Definitions and codes used to classify standard Missouri River habitats in the long term pallid sturgeon and associated fish community sampling program.....	32
Appendix C. List of standard and wild gears, their corresponding codes in the database, seasons deployed, years used, and catch-per-unit-effort units for collection of Missouri River fishes for the long-term pallid sturgeon and associated fish community sampling program	33
Appendix D. Stocking locations and codes for pallid sturgeon by Recovery Priority Management Area in the Missouri River Basin.....	34
Appendix E. Juvenile and adult pallid sturgeon stocking summary for segment 1 of the Missouri River (RPMA 2).	36
Appendix G. Hatchery names, locations, and abbreviations.	39

Introduction

The U.S. Fish and Wildlife Service (USFWS) listed the pallid sturgeon *Scaphirhynchus albus* as endangered in 1990. In response to the listing, the USFWS issued a Biological Opinion to the U.S. Army Corps of Engineers (COE), the main water management entity responsible for the Missouri River mainstem from Fort Peck Dam and Reservoir to its confluence with the Mississippi River. Additionally, an amendment to the 2000 Biological Opinion was issued in 2003. The Amendment listed several Reasonable and Prudent Alternatives (RPA) to address the inability of pallid sturgeon to naturally reproduce and the need to be able to detect changes in their populations and ecosystem trends.

The Pallid Sturgeon Population Assessment Program (Program) is guided by the RPA's in the 2003 Amendment to the 2000 Biological Opinion. The Program is a comprehensive monitoring plan designed to assess survival, movement, distribution, habitat use, and physical characteristics of these habitats used by wild and hatchery reared juvenile pallid sturgeon (Drobish 2008). The 2000 Biological Opinion divides the Program area into river and reservoir segments and assigns high, moderate, or low priority management action areas to these segments for pallid sturgeon (Drobish 2008). The focus of the Program is on the high priority management action segments. The Missouri River from Fort Peck Dam downstream to the headwaters of Lake Sakakawea, ND is listed as a high priority action segment.

The Program has stratified the Missouri River from Fort Peck Dam to the headwaters of Lake Sakakawea into four study segments based on biological, hydrological and fluvial geomorphological characteristics. The COE has contracted Montana Fish, Wildlife & Parks (FWP) to conduct Program sampling from Fort Peck Dam downstream to the North Dakota border, which consists of study segments 1 through 3.

The objectives of this program are as follows:

1. Document annual results and long-term trends in pallid sturgeon population abundance and geographic distribution throughout the Missouri River System.
2. Document annual results and long-term trends of habitat use of wild pallid sturgeon and hatchery stocked pallid sturgeon by season and life stage.

3. Document population structure and dynamics of pallid sturgeon in the Missouri River System.
4. Evaluate annual results and long-term trends in native target species population abundance and geographic distribution throughout the Missouri River system.
5. Document annual results and long-term trends of habitat usage of the native target species by season and life stage.
6. Document annual results and long-term trends of all non-target species population abundance and geographic distribution throughout the Missouri River system, where sample size is greater than fifty individuals.

Sampling Season and Species

The Program has two discrete seasons (sturgeon and fish community), which are primarily based on water temperatures. However, the sturgeon season is designed to sample sturgeon with gears that are temperature dependent, such as gill nets. Due to the nature of the majority of habitats in segment 1 through 3, gill nets are not an efficient gear for collecting pallid sturgeon due to debris flows and swift currents and therefore they are not used in any segment situated in Montana. Trammel nets and otter trawl are standard gears used in segments 1-4 during sturgeon season, and so far appear to be an effective way to sample pallid sturgeon.

The fish community season runs from the beginning of July till the end of October and is designed not only to monitor sturgeon, but also to monitor other native Missouri River fish populations. Both trammel nets and the otter trawl are still used, but to more effectively sample shallow water habitats < 1.2 m in depth mini fyke nets are also used as a standard gear.

In addition to pallid sturgeon, the Program is designed to monitor nine other native Missouri River species labeled “target” species. These include, shovelnose sturgeon *Scaphirhynchus platyrhynchus*, blue sucker *Cycleptus elongatus*, sauger *Sander canadensis*, sturgeon chub *Macrhybopsis gelida*, sicklefin chub *M. meeki*, speckled chub *M. aestivalis*, plains minnow *Hybognathus placitus*, western silvery minnow *H. argyritis*, and sand shiner *Notropis stramineus*. This suite of species was selected for various reasons. First, some species may have similar habitat requirements as pallid sturgeon and therefore by monitoring their populations we may gain further insight into pallid sturgeon habitat and how anthropomorphic and natural changes to the Missouri River affect native fish assemblages. Secondly, it is hypothesized that various chub species and other native fishes are an

important component of pallid sturgeon diet, and thereby monitoring pallids sturgeon diet will allow us to better describe their habitat. Thirdly, we wouldn't expect to see an immediate response in a long-lived species like pallid sturgeon when environmental conditions change from either favorable or detrimental conditions. Thus, by monitoring short-lived native fishes we may be able to correlate environmental conditions to changes in fish populations on a much shorter time interval and make inferences on how pallid sturgeon populations are being affected.

Study Area

Segment 1 of the Missouri River begins at Fort Peck Dam runs downstream to its confluence with the Milk River. This segment constitutes only 6% (11.5 river miles) of the entire 189.5 river miles downstream of Fort Peck Dam to the headwaters of Lake Sakakawea in North Dakota (Drobish, 2008). This reach of the Missouri River is characterized by an unnatural hydrograph, thermograph, sediment dynamics, and fish community due to the influence of Fort Peck Dam, which was constructed in 1940 (Bramblett and White, 2001). Segment 1 includes the Fort Peck Dredge Cuts, a deepened and widened section of river immediately below the dam created by the dredging of earth used to construct the dam. Regulated hypolimnetic water releases from Fort Peck Reservoir have changed a once turbid sandy bottom stretch of river into a cold clear cobble bottomed river. Fort Peck Reservoir has substantially reduced turbidity in the river below Fort Peck Dam when compared to its natural state (Galat et al, 2005).

Peaks in the hydrograph are related to power production and barge traffic downstream, instead of spring runoff and precipitation events (Galat et al, 2005). Many species native to this stretch of river such as the pallid sturgeon, sicklefin chub and sturgeon chub find the cold clear water unsuitable and are now only found farther downstream where tributaries have warmed and muddied the waters of the Missouri (Gardner and Stewart, 1987). Fish much more suited for this cold clear water such as rainbow trout *Oncorhynchus mykiss*, brown trout *Salmo trutta* and Chinook salmon *Oncorhynchus tshawytscha* have been stocked on and off from 1950 to 1990. Other nonnative species such as largemouth bass *Micropterus salmoides*, northern pike *Esox lucius*, walleye *Sander vitreus*, and yellow perch *Perca flavescens* have been stocked in the dredge cuts to increase angling opportunities. It is believed that many of these sight-feeding piscivores have out competed the native fishes in this stretch of river (Galat et al, 2005). In summary, this unique stretch of river is now vastly different from the once braided and shifting channels of the “Big Muddy” before Fort Peck Dam (Galat et al, 2005).

Methods

Sampling methods for the Pallid Sturgeon Population Assessment Program were conducted in accordance with the Standard Operating Procedures (Drobish 2008), which was established by representatives from State and Federal agencies involved with pallid sturgeon recovery on the Missouri River. For a detailed description of methodologies please see Drobish (2008). A general description of those guidelines follows.

Sampling Site Selection and Description

Montana Fish Wildlife & Parks (FWP) was contracted to sample Segment 1 from Fort Peck Dam (RM 1771.5) to the mouth of the Milk River (RM 1761), Segment 2 from the mouth of the Milk River (RM 1761) to Wolf Point (RM 1701.5) and Segment 3 from Wolf Point (RM 1701.5) to the Montana/North Dakota border (RM 1586.5). Segment 1 consists of one non-random bend at river mile 1766. This bend was selected as a reference study bend to be sampled each year to facilitate comparisons of the most highly altered area of the Missouri River in RPMA 2 (segment 1) to downstream areas (segments 2 through 4). By comparing data from segment 1 with downstream segments, a better understanding of how Fort Peck Dam influences the fish communities of the Missouri River might be attained.

During 2007 segment 1 was sampled on May 5 during the sturgeon season and September 24 to 26 during the fish community season. Two gears, trammel net and otter trawl were considered standard gears during both the sturgeon and fish community seasons, while mini fyke nets were utilized only during the fish community season.

The Population Assessment Team developed a standard set of habitat classifications for the Missouri River (Appendix B) which consists of three distinct macrohabitats found in every bend, a main channel crossover (CHXO), main channel outside bend (OSB), and main channel inside bend (ISB). Each sampling bend was comprised of these three main macrohabitats. Nine additional macrohabitats were identified that may or may not be present in every bend: large tributary mouths (TRML), small tributary mouths (TRMS), confluence areas (CONF), large and small secondary connected channels (SCCL& SCCS), deranged channels (DRNG), braided channels (BRAD), dendritic channels (DEND) and non-

connected secondary channel (SCN). For the reference bend in segment 1, five macrohabitats were sampled, CHXO, OSB, ISB, SCCL and SCCS.

Mesohabitats were established to further define macrohabitats. Mesohabitats include bars (BARS), pools (POOL), channel border (CHNB), thalweg (TLWG) and island tip (ITIP). Channel borders are situated in areas between the deepest portions of the river up to a depth of 1.2 m. Bars are considered shallow areas (< 1.2 m) where terrestrial and aquatic habitats merge. The thalweg is the deepest portion of the river between the two channel borders where the majority of the flow is directed. Pools are directly downstream of any feature that creates scour, thus creating a habitat of deep (> 1.2 m) slower moving water. Island tips are just downstream of bars or islands where two channels meet where the water is > 1.2 m in depth. Two mesohabitats were sampled in segment 1, CHNB and BARS.

For all analysis, the sampling unit was the river bend, where every river bend has a channel crossover, inside and outside bend. The downstream border of a river bend is the beginning of the next downstream bend's channel crossover.

Sampling Gear

For specific information pertaining to the specific habitats gears are utilized in and physical measurements taken in accordance with sampling the various gears described below, please see Drobis (2008).

Trammel Net

The standard trammel net has a length of 38.1 m, an inner mesh wall 2.4 m and two outer mesh walls 1.8 m deep. The inner mesh is made of #139 multifilament twine with a bar mesh size of 25.4 mm. The outer walls are constructed of #9 multifilament twine with a bar mesh size of 203.2 mm. The float line is a 12.7 mm diameter foam core with a lead line of 22.7 kg. Trammel nets were drifted from the bow of the boat and orientated perpendicular to the river flow for a minimum of 75 m and a maximum drift distance of 300 m.

Otter Trawl

The standard otter trawl has a length of 7.6 m, a width of 4.9 m and height of 0.9 m. The otter trawl has an inner mesh (6.35mm bar, #18 polyethylene twine) and outer mesh (38mmbar, #9 polyethylene twine) and a cod end opening of 406.4 mm. The trawl doors were made from 19.1 mm marine plywood and measured 762 mm x 381 mm. The trawl doors are used to keep the mouth of the trawl open while deployed on the riverbed. The trawl also has a 7.9 m long tickler chain attached to the bottom of the mouth of the trawl, which aids in keeping it orientated on the riverbed and protecting the mouth when snags are encountered. The otter trawl was deployed from the bow of the boat parallel to the current with two 30.5 m ropes and towed downstream slightly faster than current speed for a minimum of 75 m and a maximum distance of 300 m.

Mini Fyke Nets

The standard mini-fyke net consists of two rectangular frames 1.2 m wide and 0.6 m high and two 0.6 m tempered steel hoops. A 4.5 m long and 0.6 m high lead is connected to the first frame. The fyke net was made of 3 mm “ace” style mesh. The lead has small floats attached to the top and lead weights on the bottom. Mini-fyke nets are set with a “T” stake on shore and extend into river as perpendicular to the shoreline as possible or angled slightly downstream where higher velocities existed. Mini-fyke nets were set overnight and checked the following morning.

Data Collection and Analysis

A minimum of eight random subsamples were taken in macrohabitats were taken at the reference bend in segment 1. At least two subsamples (when possible) were taken using each gear in each macro habitat within the bend. More than two subsamples were taken in a macrohabitat for a gear when the number of discrete macrohabitats was less than four or less than four could be effectively sampled. If a pallid sturgeon was captured, we duplicated the sample in a non-random manner. No more than eight duplicates were taken and we would stop taking duplicates whenever two contiguous duplicate subsamples contain no pallid sturgeon. Although this non-random sampling, it gives us a better understanding of relative abundance and identifies habitats that pallid sturgeon may congregate in.

All fish were measured to the nearest mm. Fork length (FL) was used for pallid and shovelnose sturgeon, while other species were measured to TL with one exception, paddlefish *Polyodon spathula*, which were measured from the eye to the fork of the caudal fin. The first 25 fish of each species in each subsample were measured, after 25 they were counted.

Time was recorded at the beginning of each sample with all gears and an end time was always recorded when pulling mini fyke net sets. A global positioning satellite (GPS) position was taken at the beginning and end of all otter and beam trawls and trammel net drifts. One GPS location was taken for mini fyke net sets. All GPS locations were taken using a Garmin GPS 76 unit with Wide Area Augmentation System (WAAS) capability.

Sample depth was determined at the beginning, middle and end of each trawl and drift using a Lowrance X136 sonar unit. One depth was taken for mini fyke nets at the intersection of the frame and floatline using a wading rod.

Water temperature taken near the surface was recorded at every sample using the Lowrance X136 unit for trawls and trammel net drifts and using a hand held thermometer for mini fyke nets.

Habitat samples were collected randomly for 25% of each mesohabitat within each macrohabitat sampled. Velocities (mps) were taken at three depths in the water column for habitats > 1.2 m in depth (bottom, 0.8 of bottom depth and 0.2 of the bottom depth) using either a Current AA Price Meter and sounding reel or a Marsh-McBirney Flo Mate 2000. Velocities for shallow water habitats (< 1.2 m) were taken at the bottom and 0.6 of the bottom depth using the March-McBirney Flo Mate 2000.

Substrate was determined for habitats > 1.2 m in depth using a Hesse style substrate sampler. For habitats < 1.2 m in depth, a random hand grab was made. Substrate samples were reported as the percentage of sand, silt, gravel in each sample. In addition, the amount of cobble and organic material was qualitatively determined.

Turbidity was recorded in nephelometric turbidity units (NTU) using a LaMotte 2020 turbidity meter. Turbidity was taken at the midpoint of all samples, except mini fyke sets, where it was taken at the convergence of the rectangular frame and float line.

In addition to 25% of all mesohabitats, habitat measurements were taken whenever a pallid sturgeon was captured.

Genetic Verification

Genetic verification for pallid sturgeon or potential hybrids followed the methods outlined in Drobish (2008). Two fin pectoral fin clips ($\sim 2 \text{ cm}^2$) are taken from any pallid sturgeon of unknown origin. Fin samples are then preserved in 95% non-denatured alcohol for genetic analysis. All samples are sent to the U.S. Fish and Wildlife Service's Abernathy Fish Technology Center for analysis and archiving.

Analyses

The fundamental sampling unit for the Population Assessment Program is the river bend. Therefore, sample size was equal to the number of bends sampled. Accordingly, all catch-per-unit-effort (CPUE) estimates for each species by gear were made on a bend level and the mean bend CPUE's were averaged to obtain the segment CPUE.




Figure 1. Map of segment 1 of the Missouri River with major tributaries, common landmarks, and historic stocking locations for pallid sturgeon. Segment 1 encompasses the Missouri River from Fort Peck Dam (River Mile 1771.5) to the mouth of the Milk River (River Mile 1760.0).

Results

Pallid Sturgeon

No pallid sturgeon or pallid sturgeon x shovelnose sturgeon hybrids were collected in segment 1 of the Missouri River during 2007 (Figure 2).

Targeted Native River Species

Shovelnose Sturgeon

Sixteen shovelnose sturgeon were sampled in segment 1 during 2007, of which, 3 were sampled during the sturgeon season and 13 were sampled during the fish community season (Figure 2). The majority (N = 14), were captured in trammel nets, while the remaining 3 were sampled with the otter trawl (Figures 3 and 4). Trammel net CPUE was higher during both seasons when compared to otter trawl CPUE. Overall CPUE was highest for trammel net and otter trawl during the fish community season. No Shovelnose sturgeon were sampled using mini fyke nets in either 2006 or 2007.

Fewer shovelnose sturgeon were sampled during 2007 (N = 16) when compared to 2006 (N = 47). Trammel net CPUE was lower during 2007 for both the sturgeon (0.125 fish/ 100 m) and fish community seasons (0.816 fish/ 100 m) than during 2006, which had a CPUE of 1.116 and 0.947 fish/ 100 m for each season, respectively. Similarly, otter trawl CPUE was lower in 2007 for both the sturgeon (0.0 fish/100 m) and fish community (0.083 fish/ 100 m) seasons when compared to the 2006 sturgeon (0.180 fish/100 m) and fish community (0.304 fish/ 100 m) seasons.

Shovelnose sturgeon averaged 646 mm FL and weighed an average of 1,119.1 g during 2007. On average shovelnose were slightly larger during the fish community season (mean FL = 649 mm) than during the sturgeon season (mean FL = 637 mm). The length weight relationship for shovelnose sturgeon was similar between 2007 and 2006, with only a slight decrease in the slope of the power function for 2007 (Figure 6). The largest shovelnose sturgeon sampled in 2007 measured 798 mm FL, while the smallest measured 447 mm FL, which correlates to an age-5 fish (Steffensen and Hamel 2008). No young-of-the-year shovelnose sturgeon were sampled in segment 1 during either 2007 or 2006.

Sturgeon Chub

No sturgeon chubs were collected in segment 1 of the Missouri River during 2007 (Figure 2).

Sicklefin Chub

No sicklefin chubs were collected in segment 1 of the Missouri River during 2007 (Figure 2).

Sand Shiner

A total of 31 sand shiners were sampled during 2007, all in mini fyke nets during the fish community season (Figures 2 and 5). Mini fyke net CPUE for sand shiner was 3.38 fish / net night during 2007. No sand shiners were sampled in segment one during 2006.

Sand shiners averaged 52.4 mm TL with the largest and smallest samples measuring 65 and 37 mm TL, respectively. The 65 mm sand shiner was the largest specimen sampled in segments 1, 2, or 3 for both 2006 and 2007. Sand shiners in segment 1 are most likely no older than age 1+ (Dattilo et al. 2008), with possibly the exception of the 65 mm specimen.

Western Silvery Minnow

No western silvery minnows were sampled in segment 1 during 2007. Conversely, two western silvery minnows were sampled in 2006, both using the otter trawl during the sturgeon season.

Blue Sucker

One blue sucker was captured in segment 1 during 2007 in a trammel net during the sturgeon season (Figures 2 and 3). This fish measured 648 mm TL and weighed 2,500 g. Based on aging of blue suckers downstream of Gavins Point Dam, this fish is likely older than age 7 (Labay et al. 2008). Similarly, two blue suckers (657 and 690 mm TL) were sampled during 2006 in trammel nets during the sturgeon season, both were likely older than age 7.

Sauger

No sauger were sampled in segment 1 during 2007, while one was sampled during 2006 during the sturgeon season (Figure 2).




Figure 2. Total number of fish sampled during 2006 (Top Panel) and 2007 (Bottom Panel) in Segment 1 of the Missouri River. Note the break in the Y-axis' and the difference in scale below and above the break. Sample sizes are given.

Segment 1 Trammel Net CPUE




Figure 3. Trammel net CPUE by season for all target species and non-target species sampled in segment 1 of the Missouri River during 2006 (Top Panel) and 2007 (Bottom Panel). Target species are indicated by the bracket below the X-axis.

Segment 1 Otter Trawl CPUE




Figure 4. Otter trawl CPUE by season for all target species and non-target species sampled in segment 1 of the Missouri River during 2006 (Top Panel) and 2007 (Bottom Panel). Target species are indicated by the bracket below the X-axis.




Figure 5. Mini fyke net CPUE for all target species and non-target species sampled during the fish community season in segment 1 of the Missouri River during 2006 and 2007. Target species are indicated by the bracket below the X-axis (Note the differences in Y-axis for the two panels).




Figure 6. Length frequency histogram (Top Panel) for all shovelnose sturgeon sampled in Segment 1 during 2006 and 2007. Weight-length relationship (Bottom Panel) for all shovelnose sturgeon sampled in Segment 1 during 2006 and 2007. The two-parameter power functions and sample sizes are given.

Table 1. Presence absence of all species (by common name) collected in segments 1 through 3 in the Missouri River during 2006 and 2007. Boxes marked with an X indicate at least one specimen was sampled. Species in bold are native target species.

	Segments				Segments		
	1	2	3		1	2	3
Ascipenseridae - sturgeons				Esocidae - pikes			
Pallid sturgeon		X	X	Northern Pike		X	X
Shovelnose sturgeon	X	X	X	Osmeridae - smelts			
Polyodontidae - paddlefishes				Rainbow smelt			X
Paddlefish		X	X	Ictaluridae - catfishes			
Hiodontidae - mooneyes				Channel catfish	X	X	X
Goldeye	X	X	X	Black bullhead		X	X
Cyprinidae - carps and minnows				Yellow bullhead			X
Common Carp	X	X	X	Stonecat		X	X
Flathead chub	X	X	X	Salmonidae - trouts			
Emerald shiner	X	X	X	Rainbow trout	X	X	
Lake chub	X			Brown trout		X	
Longnose dace	X	X	X	Lake whitefish	X		X
Northern redbelly dace		X	X	Cisco	X		
Plains minnow		X	X	Gadidae - cods			
Western silvery minnow	X	X	X	Burbot		X	X
Brassy minnow		X		Gasterosteidae - sticklebacks			
Sicklefin chub		X	X	Brook stickleback		X	X
Sturgeon chub		X	X	Centrarchidae - sunfishes			
Sand shiner	X	X	X	Green sunfish			X
Spottail shiner	X	X	X	Pumkinseed		X	X
Fathead minnow	X	X	X	White crappie		X	X
Catostomidae-suckers				Percidae - perches			
Bigmouth buffalo		X	X	Iowa darter		X	
Smallmouth buffalo		X	X	Yellow perch	X	X	
Blue sucker	X	X	X	Sauger	X	X	X
River carpsucker	X	X	X	Walleye		X	X
White sucker	X	X	X	Sciaenidae - drums			
Longnose sucker	X	X	X	Freshwater drum			X
Shorthead redhorse	X	X	X				

Missouri River Fish Community

A total of 12,362 fish consisting of 14 species were sampled in segment 1 during 2007. White suckers *Catostomus commersoni* were the most common species sampled making up 82% of the catch (N = 10,118). The remaining 13 species in order of their abundance were, fathead minnows *Pimephales promelas* (N = 1,171), emerald shiners *Notropis atherinoides* (N = 491), longnose suckers *Catostomus catostomus* (N = 490), sand shiners (N = 31), longnose dace *Rhinichthys cataractae* (N = 26), shovelnose sturgeon (N = 16), flathead chubs *Platygobio gracilis* (N = 7), shorthead redhorse *Moxostoma macrolepidotum* (N = 4), river carpsucker *Carpionodes carpio* (N = 4), yellow perch *Perca flavescens* (N = 1), spottail shiner *Notropis hudsonius* (N = 1), common carp *Cyprinus carpio* (N = 1), and blue sucker (N = 1).

Of the total catch, only 10 fish were sampled during the sturgeon season, while mini fyke nets sampled over 99% of the total catch. During 2007, 7 species were collected that were not sampled during 2006, including four native fishes (fathead chub, river carpsucker, shorthead redhorse, and sand shiner) and three nonnative species (common carp, spottail shiner and yellow perch). Four species that were sampled in 2006 were not found in the 2007 sampling, including the native goldeye and the nonnative rainbow trout *Oncorhynchus mykiss*, lake whitefish *Coregonus clupeaformis* and ciscoe *C. artedi*.

Discussion

Segment 1 of the Missouri River is a highly altered segment due to the proximity of Fort Peck Dam. Fort Peck Dam, a hypolimnetic withdraw structure, is located approximately 5 river miles upstream of segment 1, which creates cold summer water temperatures and low turbidity. During the sturgeon season of 2007, water temperatures averaged 8.2 C° and water turbidity averaged 5 NTU's. During the fish community season, temperature increased to an average of 11.7 C°, while turbidity decreased to 3.5 NTU's. In comparison, segment 2, downstream of segment 1 had an average water temperature and turbidity of 13 C° and 1,147 NTU's during the sturgeon season and an average water temperature of 15.6 C° and turbidity of 23.2 NTU's during the fish community season. Similarly, the benthic substrate of segment 1 is noticeably different than the substrates of downstream segments, with segment 1 being composed of more gravel and coble.

Even with the highly altered conditions of segment 1, many native species are still occupying the habitats of this segment. Of the thirteen species collected, only three were not native to this section of the Missouri River (common carp, yellow perch and spottail shiner). Although no small (< 447 mm FL) shovelnose sturgeon were captured in segment 1, the trammel net CPUE for shovelnose sturgeon was similar during the fish community in segment 1 (0.816 fish/ 100m) season to that of segment 2 (0.839 fish/ 100 m). This area seems to be an adult rearing area for shovelnose sturgeon, with little or no juvenile rearing taking place. Similarly, this section of the river and the adjacent dredge cuts are known to be occupied by a relatively large number of shovelnose sturgeon during the winter months (Dave Fuller, Montana Fish, Wildlife and Parks, personal communication).

Noteworthy changes in the relative abundance of small fishes occurred in mini fyke net sampling during 2007 when compared to 2006 (Figure 5). All species collected in mini fyke nets during 2007 had higher CPUE than during 2006, with the exception of rainbow trout (*Oncorhynchus mykiss*), which were not sampled in 2007. For example, white sucker and fathead minnow CPUE averaged 1,2601 and 143.4 fish/ net night, a noticeable increase from 2006 where these species had CPUE of 0.6 and 1.3 fish/ net night, respectively. Conversely to the small sized fishes, CPUE for both trammel net and otter trawl were mostly lower for large fish during 2007 when compared to 2006 (Figures 3 and 4).

Although it is unknown why comparatively large changes in the relative abundance of fishes occurred in segment 1 between 2007 and 2006, one parameter stands out between the two years. The Milk River, which enters the Missouri River approximately 4.5 river miles downstream of segment 1 had discharges that were near 5,000 cfs during June of 2007, while during the same period in 2006 flows were remained for the most part below 200 cfs. This increase in flow from the Milk River may have caused more pre-spawning fish to move to the upper reaches of the Missouri River and Milk River during 2007 than 2006. However, no significant differences were observed in discharge between 2006 and 2007 for the Missouri River directly downstream of Fort Peck Dam and above the confluence of the Milk River. It seems unlikely that larvae fish from the Milk River would end up rearing in segment 1, due to it being upstream of the confluence of the two rivers.

No pallid sturgeon were sampled in segment 1 during 2007 or 2006, despite continuous stocking from 2004 to 2007 in the area of the Milk and Missouri Rivers confluence. Specifically during 2007, 639 yearling and 7,256 fingerlings were stocked at the School Trust stocking site on the Missouri River, which lies approximately 2 river miles downstream of segment 1. Data have indicated that at least some stocked pallid sturgeon from all other stockings sites in segments 2 and 3 do move upstream. This behavior has not been seen to date with the school trust or mouth of Milk River stocking locations.

In two years of sampling we have sampled at total of 22, 38 and 38 species in segments 1, 2 and 3 of the Missouri River, respectively (Table 1). Of those, 6 species in segment 1, 10 in segment 2 and 11 in segment 3 are not native to the Missouri River. Nonnatives sampled in segment 1 are more characteristic of cool clear water than those found in segments downstream.

Acknowledgments

The U.S. Army Corps of Engineers provided funding for this project. We'd like to thank Mark Drobish for continuing to support us in all aspects. John Hunziker, Bob Lipscomb and Landon Johnson assisted in both the field and shop throughout the year. Mike Ruggles took care many details in the office and beyond while we were busy playing on the river. We'd like to thank the entire Flow Modification Crew for their help in the field and office. Thanks to Pat Braaten of the U.S. Geological Survey for answering any type of question we may have pertaining to the Missouri River and its fishes. Thanks to Ryan Wilson and Steve Krentz of the U.S. Fish and Wildlife Service for all the collaboration between our offices.

References

- Dattilo, J. E., R. R. Dirnberger, P. T. Horner, D. J. Niswonger, M. L. Miller and V. H. Travinchek. 2008. Three Year Summary Age and Growth Report For Sand Shiner (*Notropis stramineus*). Pallid Sturgeon Population Assessment Project and Associated Fish Community Monitoring for the Missouri River. Missouri Department of Conservation. Chillicothe, MO.
- Drobish, M.R. (editor), 2008. Missouri River Standard Operating Procedures and Data Collection, Volume 1.3. U.S. Army Corps of Engineers, Omaha District, Yankton, S.D.
- Galat, D.L., C.R. Berry Jr., E.J. Peters and R.G. White. 2005. Missouri River. Pages 427-480 in A.C. Benke and C.E. Cushing (editors). Rivers of North America, Elsevier, Oxford.
- Gardner, W.M. and P.A. Stewart. 1987. The Fishery of the Lower Missouri River. Federal Aid to Fish and Wildlife Restoration Project FW-2-R Job I-b. Montana Fish, Wildlife and Parks. Helena, Montana.
- Labay, S., J. Kral and S. Stukel. 2008. Three Year Summary Age and Growth Report For Blue Sucker. Pallid Sturgeon Population Assessment Project and Associated Fish Community Monitoring for the Missouri River. South Dakota Department of Game, Fish and Parks. Yankton, SD.
- Pierce, C. L., C. S. Guy, P. J. Braaten, and M.A. Pegg. 2004. Fish growth, mortality, recruitment, condition, and size structure. Volume 4. Population structure and habitat use of benthic fishes along the Missouri and lower Yellowstone Rivers. U.S. Geological Survey, Cooperative Research Units, Iowa State University, Ames Iowa.
- Steffensen, K. and M. Hamel. 2008. Four Year Summary Age and Growth Report For Shovelnose Sturgeon. Pallid Sturgeon Population Assessment Project and Associated Fish Community Monitoring for the Missouri River. Nebraska Game and Parks Commission. Lincoln, NE.

APPENDICES

Appendix A. Phylogenetic list of Missouri River fishes with corresponding letter codes used in the long-term pallid sturgeon and associated fish community sampling program. The phylogeny follows that used by the American Fisheries Society, Common and Scientific Names of Fishes from the United States and Canada, 5th edition. Asterisks and bold type denote targeted native Missouri River species.

Scientific name	Common name	Letter Code
CLASS CEPHALASPIDOMORPHI-LAMPREYS		
ORDER PETROMYZONTIFORMES		
Petromyzontidae – lampreys		
<i>Ichthyomyzon castaneus</i>	Chestnut lamprey	CNLP
<i>Ichthyomyzon fossor</i>	Northern brook lamprey	NBLP
<i>Ichthyomyzon unicuspis</i>	Silver lamprey	SVLP
<i>Ichthyomyzon gagei</i>	Southern brook lamprey	SBLR
Petromyzontidae	Unidentified lamprey	ULY
Petromyzontidae larvae	Unidentified larval lamprey	LVLP
CLASS OSTEICHTHYES – BONY FISHES		
ORDER ACIPENSERIFORMES		
Acipenseridae – sturgeons		
<i>Acipenser fulvescens</i>	Lake sturgeon	LKSG
<i>Scaphirhynchus</i> spp.	Unidentified Scaphirhynchus	USG
<i>Scaphirhynchus albus</i>	Pallid sturgeon	PDSG*
<i>Scaphirhynchus platyrhynchus</i>	Shovelnose sturgeon	SNSG*
<i>S. albus</i> X <i>S. platyrhynchus</i>	Pallid-shovelnose hybrid	SNPD
Polyodontidae – paddlefishes		
<i>Polyodon spathula</i>	Paddlefish	PDFH
ORDER LEPISOSTEIFORMES		
Lepisosteidae – gars		
<i>Lepisosteus oculatus</i>	Spotted gar	STGR
<i>Lepisosteus osseus</i>	Longnose gar	LNGR
<i>Lepisosteus platostomus</i>	Shortnose gar	SNGR
ORDER AMMIFORMES		
Amiidae – bowfins		
<i>Amia calva</i>	Bowfin	BWFN
ORDER OSTEOGLOSSIFORMES		
Hiodontidae – mooneyes		
<i>Hiodon alosoides</i>	Goldeye	GDEY
<i>Hiodon tergisus</i>	Mooneye	MNEY
ORDER ANGUILLIFORMES		
Anguillidae – freshwater eels		
<i>Anguilla rostrata</i>	American eel	AMEL

Appendix A. (continued).

Scientific name	Common name	Letter Code
ORDER CLUPEIFORMES		
Clupeidae – herrings		
<i>Alosa alabame</i>	Alabama shad	ALSD
<i>Alosa chrysochloris</i>	Skipjack herring	SJHR
<i>Alosa pseudoharengus</i>	Alewife	ALWF
<i>Dorosoma cepedianum</i>	Gizzard shad	GZSD
<i>Dorosoma petenense</i>	Threadfin shad	TFSD
<i>D. cepedianum X D. petenense</i>	Gizzard-threadfin shad hybrid	GSTS
ORDER CYPRINIFORMES		
Cyprinidae – carps and minnows		
<i>Campostoma anomalum</i>	Central stoneroller	CLSR
<i>Campostoma oligolepis</i>	Largescale stoneroller	LSSR
<i>Carassus auratus</i>	Goldfish	GDFH
<i>Carassus auratus X Cyprinus carpio</i>	Goldfish-Common carp hybrid	GFCC
<i>Couesius plumbens</i>	Lake chub	LKCB
<i>Ctenopharyngodon idella</i>	Grass carp	GSCP
<i>Cyprinella lutrensis</i>	Red shiner	RDSN
<i>Cyprinella spiloptera</i>	Spotfin shiner	SFSN
<i>Cyprinus carpio</i>	Common carp	CARP
<i>Erimystax x-punctatus</i>	Gravel chub	GVCB
<i>Hybognathus argyritis</i>	Western slivery minnow	WSMN*
<i>Hybognathus hankinsoni</i>	Brassy minnow	BSMN
<i>Hybognathus nuchalis</i>	Mississippi silvery minnow	SVMW
<i>Hybognathus placitus</i>	Plains minnow	PNMW*
<i>Hybognathus spp.</i>	Unidentified Hybognathus	HBNS*
<i>Hypophthalmichthys molitrix</i>	Silver carp	SVCP
<i>Hypophthalmichthys nobilis</i>	Bighead carp	BHCP
<i>Luxilus chrysocephalus</i>	Striped shiner	SPSN
<i>Luxilus cornutus</i>	Common shiner	CMSN
<i>Luxilus zonatus</i>	Bleeding shiner	BDSN
<i>Lythrurus unbratilis</i>	Western redfin shiner	WRFS
<i>Macrhybopsis aestivalis</i>	Speckled chub	SKCB*
<i>Macrhybopsis gelida</i>	Sturgeon chub	SGCB*
<i>Macrhybopsis meeki</i>	Sicklefin chub	SFCB*
<i>Macrhybopsis storeriana</i>	Silver chub	SVCB
<i>M. aestivalis X M. gelida</i>	Speckled-Sturgeon chub hybrid	SPST
<i>M. gelida X M. meeki</i>	Sturgeon-Sicklefin chub hybrid	SCSC
<i>Macrhybopsis spp.</i>	Unidentified chub	UHY
<i>Margariscus margarita</i>	Pearl dace	PLDC
<i>Mylocheilus caurinus</i>	Peamouth	PEMT
<i>Nocomis biguttatus</i>	Hornyhead chub	HHCB
<i>Notemigonus crysoleucas</i>	Golden shiner	GDSN
<i>Notropis atherinoides</i>	Emerald shiner	ERSN
<i>Notropis blennioides</i>	River shiner	RVSN
<i>Notropis boops</i>	Bigeye shiner	BESN
<i>Notropis buechanani</i>	Ghost shiner	GTSN
<i>Notropis dorsalis</i>	Bigmouth shiner	BMSN
<i>Notropis greeniei</i>	Wedgespot shiner	WSSN

Appendix A. (continued).

Scientific name	Common name	Letter Code
Cyprinidae – carps and minnows		
<i>Notropis heterolepsis</i>	Blacknose shiner	BNSN
<i>Notropis hudsonius</i>	Spottail shiner	STSN
<i>Notropis nubilus</i>	Ozark minnow	OZMW
<i>Notropis rubellus</i>	Rosyface shiner	RYSN
<i>Notropis shumardi</i>	Silverband shiner	SBSN
<i>Notropis stilbius</i>	Silverstripe shiner	SSPS
<i>Notropis stramineus</i>	Sand shiner	SNSN*
<i>Notropis topeka</i>	Topeka shiner	TPSN
<i>Notropis volucellus</i>	Mimic shiner	MMSN
<i>Notropis wickliffi</i>	Channel shiner	CNSN
<i>Notropis</i> spp.	Unidentified shiner	UNO
<i>Opsopoeodus emiliae</i>	Pugnose minnow	PNMW
<i>Phenacobius mirabilis</i>	Suckermouth minnow	SMMW
<i>Phoxinus eos</i>	Northern redbelly dace	NRBD
<i>Phoxinus erythrogaster</i>	Southern redbelly dace	SRBD
<i>Phoxinus neogaeus</i>	Finescale dace	FSDC
<i>Pimephales notatus</i>	Bluntnose minnow	BNMW
<i>Pimephales promelas</i>	Fathead minnow	FHMW
<i>Pimephales vigilas</i>	Bullhead minnow	BHMW
<i>Platygobio gracilis</i>	Flathead chub	FHCB
<i>P. gracilis</i> X <i>M. meeki</i>	Flathead-sicklefin chub hybrid	FCSC
<i>Rhinichthys atratulus</i>	Blacknose dace	BNDC
<i>Rhinichthys cataractae</i>	Longnose dace	LNDC
<i>Richardsonius balteatus</i>	Redside shiner	RDSS
<i>Scardinius erythrophthalmus</i>	Rudd	RUDD
<i>Semotilus atromaculatus</i>	Creek chub	CKCB
	Unidentified Cyprinidae	UCY
	Unidentified Asian Carp	UAC
Catostomidae - suckers		
<i>Carpionodes carpio</i>	River carpsucker	RVCS
<i>Carpionodes cyprinus</i>	Quillback	QLBK
<i>Carpionodes velifer</i>	Highfin carpsucker	HFCS
<i>Carpionodes</i> spp.	Unidentified <i>Carpionodes</i>	UCS
<i>Catostomus catostomus</i>	Longnose sucker	LNSK
<i>Catostomus commersoni</i>	White sucker	WTSK
<i>Catostomus platyrhincus</i>	Mountain sucker	MTSK
<i>Catostomus</i> spp.	Unidentified <i>Catostomus</i> spp.	UCA
<i>Cycleptus elongates</i>	Blue sucker	BUSK*
<i>Hypentelium nigricans</i>	Northern hog sucker	NHSK
<i>Ictiobus bubalus</i>	Smallmouth buffalo	SMBF
<i>Ictiobus cyprinellus</i>	Bigmouth buffalo	BMBF
<i>Ictiobus niger</i>	Black buffalo	BKBF
<i>Ictiobus</i> spp.	Unidentified buffalo	UBF
<i>Minytrema melanops</i>	Spotted sucker	SPSK
<i>Moxostoma anisurum</i>	Silver redhorse	SVRH
<i>Moxostoma carinatum</i>	River redhorse	RVRH
<i>Moxostoma duquesnei</i>	Black redhorse	BKRH
<i>Moxostoma erythrurum</i>	Golden redhorse	GDRH
<i>Moxostoma macrolepidotum</i>	Shorthead redhorse	SHRH
<i>Moxostoma</i> spp.	Unidentified redhorse	URH

Appendix A. (continued).

Scientific name	Common name	Letter Code
Catostomidae - suckers	Unidentified Catostomidae	UCT
ORDER SILURIFORMES		
Ictaluridae – bullhead catfishes		
<i>Ameiurus melas</i>	Black bullhead	BKBH
<i>Ameiurus natalis</i>	Yellow bullhead	YLBH
<i>Ameiurusnebulosus</i>	Brown bullhead	BRBH
<i>Ameiurus</i> spp.	Unidentified bullhead	UBH
<i>Ictalurus furcatus</i>	Blue catfish	BLCF
<i>Ictalurus punctatus</i>	Channel catfish	CNCF
<i>I. furcatus</i> X <i>I. punctatus</i>	Blue-channel catfish hybrid	BCCC
<i>Ictalurus</i> spp.	Unidentified <i>Ictalurus</i> spp.	UCF
<i>Noturus exilis</i>	Slender madtom	SDMT
<i>Noturus flavus</i>	Stonecat	STCT
<i>Noturus gyrinus</i>	Tadpole madtom	TPMT
<i>Noturus nocturnes</i>	Freckled madtom	FKMT
<i>Pylodictis olivaris</i>	Flathead catfish	FHCF
ORDER SALMONIFORMES		
Esocidae - pikes		
<i>Esox americanus vermiculatus</i>	Grass pickerel	GSPK
<i>Esox lucius</i>	Northern pike	NTPK
<i>Esox masquinongy</i>	Muskellunge	MSKG
<i>E. lucius</i> X <i>E. masquinongy</i>	Tiger Muskellunge	TGMG
Umbridae - mudminnows		
<i>Umbra limi</i>	Central mudminnow	MDMN
Osmeridae - smelts		
<i>Osmerus mordax</i>	Rainbow smelt	RBST
Salmonidae - trouts		
<i>Coregonus artedi</i>	Lake herring or cisco	CSCO
<i>Coregonus clupeaformis</i>	Lake whitefish	LKWF
<i>Oncorhynchus aguabonita</i>	Golden trout	GDTT
<i>Oncorhynchus clarki</i>	Cutthroat trout	CTTT
<i>Oncorhynchus kisutch</i>	Coho salmon	CHSM
<i>Oncorhynchus mykiss</i>	Rainbow trout	RBTT
<i>Oncorhynchus nerka</i>	Sockeye salmon	SESM
<i>Oncorhynchus tshawytscha</i>	Chinook salmon	CNSM
<i>Prosopium cylindraceum</i>	Bonniville cisco	BVSC
<i>Prosopium williamsoni</i>	Mountain whitefish	MTWF
<i>Salmo trutta</i>	Brown trout	BNTT
<i>Salvelinus fontinalis</i>	Brook trout	BKTT
<i>Salvelinus namaycush</i>	Lake trout	LKTT
<i>Thymallus arcticus</i>	Arctic grayling	AMGL

Appendix A. (continued).

Scientific name	Common name	Letter Code
	ORDER PERCOPSIFORMES	
	Percopsidae – trout-perches	
<i>Percopsis omiscomaycus</i>	Trout-perch	TTPH
	ORDER GADIFORMES	
	Gadidae - cods	
<i>Lota lota</i>	Burbot	BRBT
	ORDER ATHERINIFORMES	
	Cyprinodontidae - killifishes	
<i>Fundulus catenatus</i>	Northern studfish	NTSF
<i>Fundulus daphanus</i>	Banded killifish	BDKF
<i>Fundulus notatus</i>	Blackstripe topminnow	BSTM
<i>Fundulus olivaceus</i>	Blackspotted topminnow	BPTM
<i>Fundulus sciadicus</i>	Plains topminnow	PTMW
<i>Fundulus zebrinus</i>	Plains killifish	PKLF
	Poeciliidae - livebearers	
<i>Gambusia affinis</i>	Western mosquitofish	MQTF
	Atherinidae - silversides	
<i>Labidesthes sicculus</i>	Brook silverside	BKSS
	ORDER GASTEROSTEIFORMES	
	Gasterosteidae - sticklebacks	
<i>Culea inconstans</i>	Brook stickleback	BKSB
	ORDER SCORPAENIFORMES	
	Cottidae - sculpins	
<i>Cottus bairdi</i>	Mottled sculpin	MDSP
<i>Cottus carolinae</i>	Banded sculpin	BDSP
	ORDER PERCIFORMES	
	Percichthyidae – temperate basses	
<i>Morone Americana</i>	White perch	WTPH
<i>Morone chrysops</i>	White bass	WTBS
<i>Morone mississippiensis</i>	Yellow bass	YWBS
<i>Morone saxatilis</i>	Striped bass	SDBS
<i>M. saxatilis X M. chrysops</i>	Striped-white bass hybrid	SBWB
	Centrarchidae - sunfishes	
<i>Ambloplites rupestris</i>	Rock bass	RKBS
<i>Archoplites interruptus</i>	Sacramento perch	SOPH
<i>Lepomis cyanellus</i>	Green sunfish	GNSF
<i>Lepomis gibbosus</i>	Pumpkinseed	PNSD
<i>Lepomis gulosus</i>	Warmouth	WRMH
<i>Lepomis humilis</i>	Orangespotted sunfish	OSSF
<i>Lepomis macrochirus</i>	Bluegill	BLGL
<i>Lepomis magalotis</i>	Longear sunfish	LESF
<i>Lepomis microlophus</i>	Redear sunfish	RESF
<i>L. cyanellus X L. macrochirus</i>	Green sunfish-bluegill hybrid	GSBG

Appendix A. (continued).

Scientific name	Common name	Letter Code
Centrarchidae - sunfishes		
<i>L. cyanellus</i> X <i>L. humilis</i>	Green-orangespotted sunfish hybrid	GSOS
<i>L. macrochirus</i> X <i>L. microlophus</i>	Bluegill-redear sunfish hybrid	BGRE
<i>Lepomis</i> spp.	Unidentified <i>Lepomis</i>	ULP
<i>Micropterus dolomieu</i>	Smallmouth bass	SMBS
<i>Micropterus punctatus</i>	Spotted sunfish	STBS
<i>Micropterus salmoides</i>	Largemouth bass	LMBS
<i>Micropterus</i> spp.	Unidentified <i>Micropterus</i> spp.	UMC
<i>Pomoxis annularis</i>	White crappie	WTCP
<i>Pomoxis nigromaculatus</i>	Black crappie	BKCP
<i>Pomoxis</i> spp.	Unidentified crappie	UCP
<i>P. annularis</i> X <i>P. nigromaculatus</i>	White-black crappie hybrid	WCBC
Centrarchidae	Unidentified centrarchid	UCN
Percidae - perches		
<i>Ammocrypta asprella</i>	Crystal darter	CLDR
<i>Etheostoma blennioides</i>	Greenside darter	GSDR
<i>Etheostoma caeruleum</i>	Rainbow darter	RBDR
<i>Etheostoma exile</i>	Iowa darter	IODR
<i>Etheostoma flabellare</i>	Fantail darter	FTDR
<i>Etheostoma gracile</i>	Slough darter	SLDR
<i>Etheostoma microperca</i>	Least darter	LTDR
<i>Etheostoma nigrum</i>	Johnny darter	JYDR
<i>Etheostoma punctulatum</i>	Stippled darter	STPD
<i>Etheostoma spectabile</i>	Orangethroated darter	OTDR
<i>Etheostoma tetrazonum</i>	Missouri saddled darter	MSDR
<i>Etheostoma zonale</i>	Banded darter	BDDR
<i>Etheostoma</i> spp.	Unidentified <i>Etheostoma</i> spp.	UET
<i>Perca flavescens</i>	Yellow perch	YWPH
<i>Percina caproides</i>	Logperch	LGPH
<i>Percina cymatotaenia</i>	Bluestripe darter	BTDR
<i>Percina evides</i>	Gilt darter	GLDR
<i>Percina maculate</i>	Blackside darter	BSDR
<i>Percina phoxocephala</i>	Slenderhead darter	SHDR
<i>Percina shumardi</i>	River darter	RRDR
<i>Percina</i> spp.	Unidentified <i>Percina</i> spp.	UPN
	Unidentified darter	UDR
<i>Sander canadense</i>	Sauger	SGER*
<i>Sander vitreus</i>	Walleye	WLEY
<i>S. canadense</i> X <i>S. vitreus</i>	Sauger-walleye hybrid/Saugeye	SGWE
<i>Sander</i> spp.	Unidentified <i>Sander</i> (formerly <i>Stizostedion</i>) spp.	UST
	Unidentified Percidae	UPC
Sciaenidae - drums		
<i>Aplodinotus grunniens</i>	Freshwater drum	FWDM
NON-TAXONOMIC CATEGORIES		
	Age-0/Young-of-year fish	YOYF
	Lab fish for identification	LAB
	No fish caught	NFSH
	Unidentified larval fish	LVFS
	Unidentified	UNID
	Net Malfunction (Did Not Fish)	NDNF

Appendix B. Definitions and codes used to classify standard Missouri River habitats in the long-term pallid sturgeon and associated fish community sampling program. Three habitat scales were used in the hierarchical habitat classification system: Macrohabitats, Mesohabitats, and Microhabitats.

Habitat	Scale	Definition	Code
Braided channel	Macro	An area of the river that contains multiple smaller channels and is lacking a readily identifiable main channel (typically associated with unchannelized sections)	BRAD
Main channel cross over	Macro	The inflection point of the thalweg where the thalweg crosses from one concave side of the river to the other concave side of the river, (i.e., transition zone from one-bend to the next bend). The upstream CHXO for a respective bend is the one sampled.	CHXO
Tributary confluence	Macro	Area immediately downstream, extending up to one bend in length, from a junction of a large tributary and the main river where this tributary has influence on the physical features of the main river	CONF
Dendric	Macro	An area of the river where the river transitions from meandering or braided channel to more of a treelike pattern with multiple channels (typically associated with unchannelized sections)	DEND
Deranged	Macro	An area of the river where the river transitions from a series of multiple channels into a meandering or braided channel (typically associated with unchannelized sections)	DRNG
Main channel inside bend	Macro	The convex side of a river bend	ISB
Main channel outside bend	Macro	The concave side of a river bend	OSB
Secondary channel-connected large	Macro	A side channel, open on upstream and downstream ends, with less flow than the main channel, large indicates this habitat can be sampled with trammel nets and trawls based on width and/or depths > 1.2 m	SCCL
Secondary channel-connected small	Macro	A side channel, open on upstream and downstream ends, with less flow than the main channel, small indicates this habitat cannot be sampled with trammel nets and trawls based on width and/or on depths < 1.2 m	SCCS
Secondary channel-non-connected	Macro	A side channel that is blocked at one end	SCCN
Tributary	Macro	Any river or stream flowing in the Missouri River	TRIB
Tributary large mouth	Macro	Mouth of entering tributary whose mean annual discharge is > 20 m ³ /s, and the sample area extends 300 m into the tributary	TRML
Tributary small mouth	Macro	Mouth of entering tributary whose mean annual discharge is < 20 m ³ /s, mouth width is > 6 m wide and the sample area extends 300 m into the tributary	TRMS
Wild	Macro	All habitats not covered in the previous habitat descriptions	WILD
Bars	Meso	Sandbar or shallow bank-line areas with depth < 1.2 m	BARS
Pools	Meso	Areas immediately downstream from sandbars, dikes, snags, or other obstructions with a formed scour hole > 1.2 m	POOL
Channel border	Meso	Area in the channelized river between the toe and the thalweg, area in the unchannelized river between the toe and the maximum depth	CHNB
Dam Tailwaters	Meso	Area below dam	DTWT
Thalweg	Meso	Main channel between the channel borders conveying the majority of the flow	TLWG
Island tip	Meso	Area immediately downstream of a bar or island where two channels converge with water depths > 1.2 m	ITIP

Appendix C. List of standard and wild gears (type), their corresponding codes in the database, seasons deployed (Fall-Spring, Summer, or all), years used, and catch-per-unit-effort units for collection of Missouri River fishes in segment 1 for the long-term pallid sturgeon and associated fish community sampling program. Long-term monitoring began in 2006 for segment 1.

Gear	Code	Type	Season	Years	CPUE units
Gillnet – 4 meshes, small mesh set upstream	GN14	Wild	Sturgeon	NOT USED	fish/net night
Gillnet – 4 meshes, large mesh set upstream	GN41	Wild	Sturgeon	NOT USED	fish/net night
Gillnet – 8 meshes, small mesh set upstream	GN18	Wild	Sturgeon	NOT USED	fish/net night
Gillnet – 8 meshes, large mesh set upstream	GN81	Wild	Sturgeon	NOT USED	fish/net night
Mini-fyke net	MF	Standard	Fish Comm.	2006 - Present	fish/net night
Push Trawl – 8 ft 4mm x 4mm	POT02	Evaluation	Fish Comm.	2006 - Present	fish/ 100 m trawled
Trammel net – 1 inch inner mesh	TN	Standard	All	2006 - Present	fish/100 m drift
Trot Line – Circle hooks**	TLC_	Wild	Sturgeon	2007 - Present	fish/hook night
Trot Line – Octopus hooks**	TLO_	Wild	Sturgeon	2007 - Present	fish/hook night
Trot Line – O’Shaughnessy hooks**	TLS_	Wild	Sturgeon	2007 - Present	fish/hook night
Otter trawl – 16 ft head rope	OT16	Standard	All	2006 - Present	fish/100 m trawled
Otter trawl – 16 ft SKT 4mm x 4mm HB2 MOR	OT01	Wild	Fish Comm.	NOT USED	fish/100 m trawled

* Standard only in upper Missouri River segments

** Code ends with line length in feet (1 = 105 ft, 2 = 205 ft, 3 = 305 ft, 4 = 405 ft). Hooks are placed between 5 and 10 feet apart.

Appendix D. Stocking locations and codes by Recovery Priority Management Area (RPMA) in the Missouri River Basin.

State(s)	RPMA	Site Name	Code	River	RM
MT	2	Forsyth	FOR	Yellowstone	253.2
MT	2	Cartersville	CAR	Yellowstone	235.3
MT	2	Miles City	MIC	Yellowstone	181.8
MT	2	Fallon	FAL	Yellowstone	124
MT	2	Intake	INT	Yellowstone	70
MT	2	Sidney	SID	Yellowstone	31
MT	2	Big Sky Bend	BSB	Yellowstone	17
ND	2	Fairview	FRV	Yellowstone	9
MT	2	Milk River	MLK	Milk	11.5
MT	2	Mouth of Milk	MOM	Missouri	1761.5
MT	2	Grand Champs	GRC	Missouri	1741
MT	2	Wolf Point	WFP	Missouri	1701.5
MT	2	Poplar	POP	Missouri	1649.5
MT	2	Brockton	BRK	Missouri	1678
MT	2	Culbertson	CBS	Missouri	1621
MT	2	Nohly Bridge	NOB	Missouri	1590
ND	2	Confluence	CON	Missouri	1581.5
SD/NE	3	Sunshine Bottom	SUN	Missouri	866.2
SD/NE	3	Verdel Boat Ramp	VER	Missouri	855
SD/NE	3	Standing Bear Bridge	STB	Missouri	845
SD/NE	3	Running Water	RNW	Missouri	840.1
SD/NE	4	St. Helena	STH	Missouri	799
SD/NE	4	Mullberry Bend	MUL	Missouri	775
NE/IA	4	Ponca State Park	PSP	Missouri	753
NE/IA	4	Sioux City	SIO	Missouri	732.6
NE/IA	4	Sloan	SLN	Missouri	709
NE/IA	4	Decatur	DCT	Missouri	691
NE/IA	4	Boyer Chute	BYC	Missouri	637.4
NE/IA	4	Bellevue	BEL	Missouri	601.4
NE/IA	4	Rulo	RLO	Missouri	497.9
NE/MO/KS	4	Kansas River	KSR	Missouri	367.5
NE	4	Platte River	PLR	Platte	5
KA/MO	4	Leavenworth	LVW	Missouri	397
MO	4	Parkville	PKV	Missouri	377.5
MO	4	Kansas City	KAC	Missouri	342

State(s)	RPMA	Site Name	Code	River	RM
MO	4	Miami	MIA	Missouri	262.8
MO	4	Grand River	GDR	Missouri	250
MO	4	Boonville	BOO	Missouri	195.1
MO	4	Overton	OVT	Missouri	185.1
MO	4	Hartsburg	HAR	Missouri	160
MO	4	Jefferson City	JEF	Missouri	143.9
MO	4	Mokane	MOK	Missouri	124.7
MO	4	Hermann	HER	Missouri	97.6
MO	4	Washington	WAS	Missouri	68.5
MO	4	St. Charles	STC	Missouri	28.5

Appendix E. Juvenile and adult pallid sturgeon stocking summary for segment 2 of the Missouri River (RPMA 2)

Year	Stocking Site	Number Stocked	Year Class	Stock Date	Age at Stocking ^a	Primary Mark	Secondary Mark
1998	Big Sky Bend	255	1997	8/11/1998	Yearling	PIT Tag	Elastomer
1998	Confluence	40	1997	8/11/1998	Yearling	PIT Tag	Elastomer
1998	Nohly Bridge	255	1997	8/11/1998	Yearling	PIT Tag	Elastomer
1998	Sidney	230	1997	8/11/1998	Yearling	PIT Tag	Elastomer
2000	Culbertson	34	1998	10/11/2000	2 yr Old	PIT Tag	
2000	Fairview	66	1998	10/11/2000	2 yr Old	PIT Tag	
2000	Sidney	66	1998	10/11/2000	2 yr Old	PIT Tag	
2000	Wolf Point	34	1998	10/11/2000	2 yr Old	PIT Tag	
2000	Culbertson	89	1999	10/17/2000	Yearling	PIT Tag	
2000	Fairview	150	1999	10/17/2000	Yearling	PIT Tag	
2000	Sidney	149	1999	10/17/2000	Yearling	PIT Tag	
2000	Wolf Point	90	1999	10/17/2000	Yearling	PIT Tag	
2002	Culbertson	270	2001	7/18/2002	Yearling	CWT	Elastomer
2002	Fairview	270	2001	7/18/2002	Yearling	CWT	Elastomer
2002	Intake	199	2001	7/18/2002	Yearling	CWT	Elastomer
2002	Sidney	271	2001	7/18/2002	Yearling	CWT	Elastomer
2002	Wolf Point	269	2001	7/18/2002	Yearling	CWT	Elastomer
2002	Culbertson	317	2001	7/26/2002	Yearling	PIT Tag	
2002	Fairview	360	2001	7/26/2002	Yearling	PIT Tag	
2002	Intake	97	2001	7/26/2002	Yearling	PIT Tag	
2002	Sidney	427	2001	7/26/2002	Yearling	PIT Tag	
2002	Wolf Point	425	2001	7/26/2002	Yearling	PIT Tag	
2002	Intake	155	2001	9/18/2002	Yearling	PIT Tag	
2003	Culbertson	1033	2002	8/7/2003	Yearling	PIT Tag	Elastomer
2003	Fairview	887	2002	8/7/2003	Yearling	PIT Tag	Elastomer
2003	Intake	1040	2002	8/7/2003	Yearling	PIT Tag	Elastomer
2003	Wolf Point	926	2002	8/7/2003	Yearling	PIT Tag	Elastomer
2004	Milk River	821	2003	4/13/2004	Yearling	Elastomer	
2004	Culbertson	523	2003	8/9/2004	Yearling	PIT Tag	Elastomer
2004	Intake	347	2003	8/9/2004	Yearling	PIT Tag	Elastomer
2004	Sidney	397	2003	8/9/2004	Yearling	PIT Tag	Elastomer
2004	Wolf Point	379	2003	8/9/2004	Yearling	PIT Tag	Elastomer
2004	Larval Drift	30000	2004	7/2/2004	Fry		
2004	Larval Drift	50000	2004	7/8/2004	Fry		
2004	Larval Drift	25000	2004	7/20/2004	Fry		
2004	Larval Drift	25000	2004	7/23/2004	Fry		
2004	Larval Drift	25000	2004	7/27/2004	Fry		
2004	Culbertson	3819	2004	9/10/2004	Fingerling	CWT	Elastomer
2004	Sidney	2991	2004	9/10/2004	Fingerling	CWT	Elastomer

Year	Stocking Site	Number Stocked	Year Class	Stock Date	Age at Stocking ^a	Primary Mark	Secondary Mark
2004	Wolf Point	4040	2004	9/10/2004	Fingerling	CWT	Elastomer
2004	Mouth of Milk	3482	2004	10/15/2004	Advanced Fingerling	CWT	Elastomer
2004	Intake	2477	2004	11/18/2004	Advanced Fingerling	CWT	Elastomer
2005	Culbertson	288	2004	4/12/2005	Yearling	CWT	Elastomer
2005	Intake	309	2004	4/12/2005	Yearling	CWT	Elastomer
2005	Wolf Point	271	2004	4/12/2005	Yearling	CWT	Elastomer
2005	Intake	175	2004	8/19/2005	Yearling	PIT Tag	Elastomer
2005	Brockton	229	2005	10/5/2005	Advanced Fingerling	CWT	Elastomer
2005	Culbertson	226	2005	10/5/2005	Advanced Fingerling	CWT	Elastomer
2005	Intake	456	2005	10/5/2005	Advanced Fingerling	CWT	Elastomer
2005	Milk River	232	2005	10/5/2005	Advanced Fingerling	CWT	Elastomer
2005	Sidney	122	2005	10/5/2005	Advanced Fingerling	CWT	Elastomer
2005	Wolf Point	611	2005	10/12/2005	Advanced Fingerling	CWT	Elastomer
2005	Brockton	371	2005	10/13/2005	Advanced fingerling		
2005	Culbertson	1736	2005	10/13/2005	Advanced Fingerling	CWT	Elastomer
2005	Culbertson	182	2005	10/13/2005	Advanced Fingerling		
2005	Intake	313	2005	10/13/2005	Advanced Fingerling		
2005	Milk River	845	2005	10/13/2005	Advanced Fingerling	CWT	Elastomer
2005	Mouth of Milk	371	2005	10/13/2005	Advanced Fingerling		
2005	Sidney	105	2005	10/13/2005	Advanced Fingerling		
2005	Wolf Point	1521	2005	10/13/2005	Advanced Fingerling	CWT	Elastomer
2005	Wolf Point	371	2005	10/13/2005	Advanced Fingerling		
2005	Culbertson	651	2005	10/19/2005	Advanced Fingerling	CWT	Elastomer
2005	Intake	2120	2005	10/19/2005	Advanced Fingerling	CWT	Elastomer
2005	Milk River	485	2005	10/19/2005	Advanced Fingerling	CWT	Elastomer
2005	Sidney	882	2005	10/19/2005	Advanced Fingerling	CWT	Elastomer
2005	Wolf Point	650	2005	10/19/2005	Advanced Fingerling	CWT	Elastomer
2006	Culbertson	235	2005	3/28/2006	Advanced Fingerling	Elastomer	
2006	Intake	327	2005	3/28/2006	Advanced Fingerling	Elastomer	
2006	Mouth of Milk	134	2005	3/28/2006	Advanced fingerling	Elastomer	
2006	Sidney	113	2005	3/28/2006	Advanced Fingerling	Elastomer	
2006	Wolf Point	232	2005	3/28/2006	Advanced Fingerling	Elastomer	
2006	Intake	970	2005	4/3/2006	Yearling	PIT Tag	Elastomer
2006	Sidney	314	2005	4/3/2006	Yearling	PIT Tag	Elastomer
2006	Culbertson	844	2005	4/5/2006	Yearling	PIT Tag	Elastomer
2006	Mouth of Milk	1007	2005	4/5/2006	Yearling	PIT Tag	Elastomer
2006	Wolf Point	866	2005	4/5/2006	Yearling	PIT Tag	Elastomer
2006	Culbertson	669	2005	5/1/2006	Yearling	PIT Tag	Scute Removed
2006	Intake	765	2005	5/1/2006	Yearling	PIT Tag	Scute Removed
2006	Mouth of Milk	650	2005	5/1/2006	Yearling	PIT Tag	Scute Removed
2006	Sidney	228	2005	5/1/2006	Yearling	PIT Tag	Scute Removed
2006	Wolf Point	653	2005	5/1/2006	Yearling	PIT Tag	Scute Removed
2006		1355	2005	5/1/2006	Yearling	PIT Tag	Scute Removed
2006	Culbertson	1544	2006	10/24/2006	Advanced Fingerling	Elastomer	
2006	Intake	1680	2006	10/24/2006	Advanced Fingerling	Elastomer	

Year	Stocking Site	Number Stocked	Year Class	Stock Date	Age at Stocking ^a	Primary Mark	Secondary Mark
2006	Mouth Milk	1117	2006	10/24/2006	Advanced Fingerling	Elastomer	
2006	Sidney	586	2006	10/24/2006	Advanced Fingerling	Elastomer	
2006	Wolf Point	1553	2006	10/24/2006	Advanced Fingerling	Elastomer	
2006	School Trust	436	2006	11/8/2006	Advanced Fingerling	Elastomer	
2007	Culbertson	651	2006	4/5/2007	Yearling	PIT Tag	Scute Removed
2007	Fallon	491	2006	4/3/2007	Yearling	PIT Tag	Scute Removed
2007	Forsyth	492	2006	4/3/2007	Yearling	PIT Tag	Scute Removed
2007	Sidney	983	2006	4/3/2007	Yearling	PIT Tag	Scute Removed
2007	School Trust	639	2006	4/5/2007	Yearling	PIT Tag	Scute Removed
2007	Wolf Point	651	2006	4/5/2007	Yearling	PIT Tag	Scute Removed
2007	Wolf Point	428285	2007	7/9/2007	Fry		
2007	Grand Champs	5558	2007	7/13/2007	Fry		
2007	Miles City	13125	2007	7/18/2007	Fry		
2007	Intake	20763	2007	8/9/2007	Fry		
2007	Miles City	13675	2007	8/9/2007	Fry		
2007	Intake	336	2007	8/27/2007	Fingerling		
2007	Miles City	336	2007	8/27/2007	Fingerling		
2007	Wolf Point	672	2007	8/27/2007	Fingerling		
2007	Forsyth	690	2007	8/31/2007	Fingerling	CWT	
2007	Intake	615	2007	8/31/2007	Fingerling	CWT	
2007	School Trust	1160	2007	9/6/2007	Fingerling	CWT	
2007	Intake	293	2007	9/12/2007	Fingerling		
2007	Miles City	293	2007	9/12/2007	Fingerling		
2007	Wolf Point	586	2007	9/12/2007	Fingerling		
2007	Culbertson	6455	2007	9/14/2007	Fingerling	Elastomer	
2007	Fallon	4827	2007	9/14/2007	Fingerling	Elastomer	
2007	Forsyth	5370	2007	9/14/2007	Fingerling	Elastomer	
2007	Intake	7812	2007	9/14/2007	Fingerling	Elastomer	
2007	School Trust	6096	2007	9/14/2007	Fingerling	Elastomer	
2007	Sidney	1934	2007	9/14/2007	Fingerling	Elastomer	
2007	Wolf Point	6455	2007	9/14/2007	Fingerling	Elastomer	

^aAge of fish when stocked: Fry, Fingerling, Yearling, 1yo, 2yo, 3yo, etc

Appendix G. Hatchery names, locations, and abbreviations.

Hatchery	State	Abbreviation
Blind Pony State Fish Hatchery	MO	BYP
Neosho National Fish Hatchery	MO	NEO
Gavins Point National Fish Hatchery	SD	GAV
Garrison Dam National Fish Hatchery	ND	GAR
Miles City State Fish Hatchery	MT	MCH
Blue Water State Fish Hatchery	MT	BLU
Bozeman Fish Technology Center	MT	BFT
Fort Peck State Fish Hatchery	MT	FPH
